

Anti-malarial Treatment Selector

Charts revised February 2019. Full information available at www.hiv-druginteractions.org

For personal use only. Not for distribution. For personal use only. Not for distribution. For personal use only. Not for distribution. For personal use only. Not for distribution.

	ATV/c	ATV/r	DRV/c	DRV/r	LPV/r	DOR	EFV	ETV	NVP	RPV	MVC	BIC/ F/TAF	DTG	EVG/c/ F/TAF	EVG/c/ F/TDF	RAL	FTC or 3TC	F/TAF	TDF	ZDV
First line and Second line Drugs																				
Amodiaquine	↑	↑	↔	↑	↑	↔	↑ ^a	↓?	↓29% ^a	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔ ^b
Artemisinin	↑	↑	↑	↑	↑	↓	↓~50%	↓↓	↓↓	↓	↓	↓ ^c	↔	↑	↑	↔	↔	↔	↔	↔
Atovaquone	↔	↔ ^d	↔	↓ ^d	↓74% ^d	↔	↓75% ^d	↑55% ^d	↓ ^d	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔ ^b
Chloroquine	↑♥	↑♥	↑	↑	↔♥	↔	↔	↔	↔	↔♥	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔
Clindamycin	↑	↑	↑	↑	↑	↔	↓	↓	↓	↔	↔	↔	↔	↑	↑	↔	↔	↔	↔	↔
Doxycycline	↔	↔	↔	↔	↔	↔	↓?	↓?	↓?	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔
Lumefantrine	↑♥	↑♥	↑	↑	↑♥	↔	↓~40%	↓	↓46%	↔♥	↔	↔	↔	↔	↑	↑	↔	↔	↔	↔
Mefloquine	↑♥	↑♥	↑	↑	↓28%♥ ↓22%	↔	↓	↓	↓	↔♥	↔	↔	↔	↑	↑	↔	↔	↔	↔	↔
Primaquine	↔	↔	↔	↔	↔	↔	↔ ^e	↔ ^e	↔ ^e	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔ ^b
Proguanil	↔	↓41% ^d	↔	↓ ^d	↓38% ^d	↔	↓44% ^d	↑↑ ^d	↓ ^d	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔
Pyrimethamine	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↑ ^f	↔	↑ ^f	↑ ^f	↔	↑ ^g	↑ ^f	↔	↔ ^b
Quinine	↑ ^h ♥	↑ ^h ♥	↑ ^h	↑ ^h	↓~50%♥	↔	↓	↓	↓	↔♥	↑	↔	↔	↑ ^h	↑ ^h	↔	↔	↔	↔	↔
Sulfadoxine	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↑ ⁱ	↔	↑ ⁱ	↑ ⁱ	↔	↑ ^j	↑ ^j	↔	↔ ^b

Colour Legend

	No clinically significant interaction expected.
	These drugs should not be coadministered.
	Potential interaction which may require a dose adjustment or close monitoring.
	Potential interaction predicted to be of weak intensity. No <i>a priori</i> dosage adjustment is recommended.

Text Legend

- ↑ Potential increased exposure of the anti-malarial drug
- ↓ Potential decreased exposure of the anti-malarial drug
- ↔ No significant effect
- ↑ Potential increased exposure of HIV drug
- ↓ Potential decreased exposure of HIV drug
- ♥ One or both drugs may cause QT and/or PR prolongation. ECG monitoring is advised if coadministered with atazanavir or lopinavir; caution is advised with rilpivirine as supratherapeutic doses of rilpivirine (75 and 300 mg once daily) were shown to prolong the QT interval.
- Numbers refer to increase or decrease in AUC as observed in drug-drug interaction studies.

Notes

- a Liver toxicity
- b Additive haematotoxicity
- c No effect on FTC or TAF is expected, but bictegravir concentrations may decrease.
- d Take with a high fat meal. Consider dose increase.
- e Increase of haemotoxic metabolites
- f FTC exposure may increase; no *a priori* dosage adjustment is recommended in patients with normal renal function.
- g FTC or 3TC exposure may increase; no *a priori* dosage adjustment is recommended in patients with normal renal function.
- h An increase in exposure would be expected based on quinine metabolism, however, two interaction studies with LPV/r have shown a decrease in quinine exposure. It is recommended to monitor for side effects and also efficacy.
- i Sulfadoxine is rarely used alone, but is usually given in combination with pyrimethamine. Pyrimethamine may increase FTC exposure, but no *a priori* dosage adjustment is recommended in patients with normal renal function.
- j Sulfadoxine is rarely used alone, but is usually given in combination with pyrimethamine. Pyrimethamine may increase FTC or 3TC exposure, but no *a priori* dosage adjustment is recommended in patients with normal renal function.