


# Prescribing in Elderly PLWH

Revised July 2019

For personal use only. Not for distribution. For personal use only. Not for distribution. For personal use only. Not for distribution.



The **Beers** and **STOPP** criteria are tools established by experts in geriatric pharmacotherapy to detect and reduce the burden of inappropriate prescribing in elderly. Inappropriate medicines include:

- Those which in elderly persons with certain diseases can lead to drug-disease interactions.
- Those associated with a higher risk of adverse drug reactions in the elderly.
- Those that predictably increase the risk of falls in the elderly.
- Those to be avoided in case of organ dysfunction.

The **START** criteria consist of evidence-based indicators of potential prescribing omission in elderly with specific medical conditions.

## References

1. Reconsidering medication appropriateness for patients late in life. Holmes HM et al. *Arch Intern Med*, 2006, 166(6): 605-9.
2. American Geriatrics Society 2019 Updated Beers Criteria for Potentially Inappropriate Medication Use in Older Adults. American Geriatrics Society Beers Criteria Update Expert Panel. *J Am Geriatr Soc*, 2019, 67(4):674-694.
3. STOPP/START criteria for potentially inappropriate prescribing in older people: version 2. O'Mahony D et al. *Age Aging*, 2015, 44(2):213-8.

# Top Ten Drug Classes to Avoid in Elderly PLWH

Produced July 2019

For personal use only. Not for distribution. For personal use only. Not for distribution. For personal use only. Not for distribution.

Drug class	Problems	Alternatives
<b>First generation antihistamines</b> e.g., Clemastine Diphenhydramine Doxylamine Hydroxyzine	Strong anticholinergic properties, risk of impaired cognition, delirium, falls, peripheral anticholinergic adverse reactions (dry mouth, constipation, blurred vision, urinary retention).	Cetirizine Desloratadine Loratadine
<b>Tricyclic antidepressants</b> e.g., Amitriptyline Clomipramine Doxepin Imipramine Trimipramine	Strong anticholinergic properties, risk of impaired cognition, delirium, falls, peripheral anticholinergic adverse reactions (dry mouth, constipation, blurred vision, urinary retention).	Citalopram Escitalopram Mirtazapine Venlafaxine
<b>Benzodiazepines</b> Long and short acting benzodiazepines e.g., Clonazepam Diazepam Midazolam Non-benzodiazepines hypnotics e.g., Zolpidem Zopiclone	Elderly are more sensitive to their effect, risk of falls, fractures, delirium, cognitive impairment, drug dependency. Use with caution, at the lowest dose, and for a short duration.	Non-pharmacological treatment of sleep disturbance/sleep hygiene
<b>Atypical antipsychotics</b> e.g., Clozapine, Olanzapine, Quetiapine	Anticholinergic adverse reactions, increased risk of stroke and mortality (all antipsychotics).	Aripiprazole Ziprasidone
<b>Urological spasmolytic agents</b> e.g., Oxybutynin, Solifenacin, Tolterodine	Strong anticholinergic properties, risk of impaired cognition, delirium, falls, peripheral anticholinergic adverse reactions (dry mouth, constipation, blurred vision, urinary retention).	Non-pharmacological treatment (pelvic floor exercises)
<b>Stimulant laxatives</b> e.g., Senna, Bisacodyl	Long-term use may cause bowel dysfunction.	Fibre Hydration Osmotic laxatives
<b>Non-steroidal anti-inflammatory drugs (NSAIDs)</b> e.g., Diclofenac, Indomethacin, Ketorolac, Naproxen	Avoid regular, long-term use of NSAIDs due to risk of gastro-intestinal bleeding, renal failure, worsening of heart failure.	Paracetamol Weak opioids
<b>Cardiac glycosides</b> Digoxin (>0.125 mg/day)	Avoid doses higher than 0.125 mg/day due to risk of toxicity.	Beta-blockers (for atrial fibrillation)
<b>Long acting sulfonylureas</b> e.g., Glyburide, Chlorpropamide	Can cause severe prolonged hypoglycaemia.	Metformin or other antidiabetic classes
<b>Cold medications</b> Most of these products contain antihistamines (e.g., diphenhydramine) and decongestants (e.g., phenylephrine, pseudoephedrine)	First generation antihistamines can cause central and peripheral anticholinergic adverse reactions (as described above). Oral decongestants can increase blood pressure.	Avoid

Refer to Beers criteria for a more exhaustive list:

American Geriatrics Society 2019 updated AGS Beers Criteria for potentially inappropriate medication use in older adults.  
2019 American Geriatrics Society Beers Criteria Update Expert Panel. *J Am Geriatr Soc*, 2019, 67(4):674-694.

# Common Prescribing Cascades to Avoid in Elderly PLWH

Produced July 2019

For personal use only. Not for distribution. For personal use only. Not for distribution. For personal use only. Not for distribution.

Initial treatment	Adverse drug reaction	Subsequent treatment
ACE inhibitors	→ Cough	→ Cough suppressant; antibiotic
Amlodipine	→ Edema	→ Diuretics
Antihypertensives	→ Dizziness	→ Prochlorperazine
Antipsychotics	→ Extrapyramidal effect	→ Antiparkinsonian agents
Beta-blockers	→ Depression	→ Antidepressants
Cholinesterase inhibitors	↗ Incontinence	→ Anticholinergics
	→ Diarrhoea	→ Bismuth subsalicylate
	↘ Rhinorrhoea	→ Diphenhydramine
Erythromycin	→ Arrhythmia	→ Antiarrhythmics
Lithium	→ Tremor	→ Propranolol
Meperidine	→ Delirium	→ Antipsychotics*
Metoclopramide	→ Extrapyramidal effect	→ Antiparkinsonian agents
NSAIDs	→ Rise in blood pressure	→ Antihypertensives
Quinolone	→ Delirium	→ Antipsychotics*
SSRI; SNRI	→ Tremor	→ Benzodiazepines
Thiazide diuretics	→ Hyperuricemia; gout	→ Allopurinol; colchicine
Tricyclic antidepressants	↗ Decreased cognition	→ Cholinesterase inhibitors*
	↘ Constipation	→ Laxatives

ACE = angiotensin-converting enzyme  
 NSAID = nonsteroidal anti-inflammatory drug  
 SNRI = serotonin-norepinephrine reuptake inhibitor  
 SSRI = selective serotonin reuptake inhibitor

\* Subsequent treatment could result in a further prescribing cascade

## References

- Caughey GE, Roughhead EE, Pratt N, Shakib S, Vitry AI, Gilbert AL. Increased risk of hip fracture in the elderly associated with prochlorperazine: is a prescribing cascade contributing?  
*Pharmacoepidemiol Drug Saf.* 2010, 19(9):977-82.
- Corrao G, Botteri E, Bagnardi V, et al. Generating signals of drug-adverse effects from prescription databases and application to the risk of arrhythmia associated with antibacterials.  
*Pharmacoepidemiol Drug Saf.* 2005, 14(1):31-40.
- Gill SS, Mamdani M, Naglie G, et al. A prescribing cascade involving cholinesterase inhibitors and anticholinergic drugs.  
*Arch Intern Med.* 2005, 165(7):808-13.
- Gurwitz JH, Avorn J, Bohn RL, et al. Initiation of antihypertensive treatment during nonsteroidal anti-inflammatory drug therapy.  
*JAMA.* 1994, 272(10):781-6.
- Liu PT, Argento VS, Skudlarska BA. Prescribing cascade in an 80-year-old Japanese immigrant. *Geriatr Gerontol Int.* 2009, 9(4):402-4.
- Nguyen PV, Spinelli C. Prescribing cascade in an elderly woman. *Can Pharm J (Ott).* 2016, 149(3):122-4.
- Onder G, Bonassi S, Abbatecola AM, et al. High prevalence of poor quality drug prescribing in older individuals: a nationwide report from the Italian Medicines Agency (AIFA).  
*J Gerontol A Biol Sci Med Sci.* 2014, 69(4):430-7.
- Rosenberg J, Rochon PA, Gill SS. Unveiling a prescribing cascade in an older man. *J Am Geriatr Soc.* 2014, 62(3):580-1.
- Vegter S, de Jong-van den Berg LT. Misdiagnosis and mistreatment of a common side-effect--angiotensin-converting enzyme inhibitor-induced cough. *Br J Clin Pharmacol.* 2010, 69(2):200-3.
- Veloso TS, Cambao MS. Migraine treatment: a chain of adverse effects. *Springerplus.* 2015, 4:409.
- Vouri SM, Chung JM, Binder EF. Successful intervention to mitigate an acetylcholinesterase inhibitor-induced rhinorrhea prescribing cascade: a case report. *J Clin Pharm Ther.* 2017, 42(3):370-1.