

# Anti-diabetic Treatment Selector

Charts reviewed June 2022. Full information available at [www.hiv-druginteractions.org](http://www.hiv-druginteractions.org)

For personal use only. Not for distribution. For personal use only. Not for distribution. For personal use only. Not for distribution. For personal use only. Not for distribution.

	ATV/c	ATV/r	DRV/c	DRV/r	LPV/r	DOR	EFV	ETV	NVP	RPV oral	FTR	MVC	BIC/F/TAF	CAB oral	CAB/RPV	DTG	EVG/c/F/TAF	EVG/c/F/TDF	RAL	FTC/TAF	FTC/TDF	TDF
<b>Sulfonylureas</b>																						
Glibenclamide	↑	↑	↑	↑	↑	↔	↓	↓	↓	↔	↑	↔	↔	↔	↔	↔	↑	↑	↔	↔	↔	↔
Gliclazide	↔	↓	↔	↓	↓	↔	↑	↑	↔	↔	↑	↔	↔	↔	↔	↔	↓	↓	↔	↔	↔	↔
Glimepiride	↔	↓	↔	↓	↓	↔	↑	↑	↔	↔	↑	↔	↔	↔	↔	↔	↓	↓	↔	↔	↔	↔
Glipizide	↔	↓	↔	↓	↓	↔	↑	↑	↔	↔	↑	↔	↔	↔	↔	↔	↓	↓	↔	↔	↔	↔
Tolbutamide	↔	↓	↔	↓	↓	↔	↑	↑	↔	↔	↔	↔	↔	↔	↔	↔	↓	↓	↔	↔	↔	↔
<b>Biguanides</b>																						
Metformin	↑ a	↔	↑ a	↔	↔	↓ 6%	↔	↔	↔	↓ 3%	↔	↔	↑ 39%	↔	↔	↑ 79% a	↑ a	↑ a	↔	↔	↔	↔
<b>Thiazolidinediones</b>																						
Pioglitazone	↑	↑	↑	↑	↑	↔	↑	↓	↓	↔	↔	↔	↔	↔	↔	↔	↑	↑	↔	↔	↔	↔
Rosiglitazone	↑ 35%	↓ 17%	↔	↓	↓	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔
<b>Meglitinides</b>																						
Nateglinide	↑	↑↓	↑	↑↓	↑↓	↔	↑↓	↑↓	↓	↔	↑	↔	↔	↔	↔	↔	↑↓	↑↓	↔	↔	↔	↔
Repaglinide	↑	↑	↑	↑	↑	↔	↑↓	↓	↓	↔	↑	↔	↔	↔	↔	↔	↑	↑	↔	↔	↔	↔
<b>GLP-1 agonists</b>																						
Dulaglutide	↔↓	↔↓	↔	↔	↔	↔	↔	↔	↔	↔↓	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔
Exenatide	↔↓ b	↔↓ b	↔	↔	↔	↔	↔	↔	↔	↔↓ c	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔
Liraglutide	↔↓ b	↔↓ b	↔	↔	↔	↔	↔	↔	↔	↔↓ c	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔
Semaglutide	↔↓ b	↔↓ b	↔	↔	↔	↔	↔	↔	↔	↔↓ c	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔
<b>DPP-4 inhibitors</b>																						
Alogliptin	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔
Linagliptin	↑ d	↑ d	↑ d	↑ d	↑ d	↔	↓	↓	↓	↔	↔	↔	↔	↔	↔	↔	↑ d	↑ d	↔	↔	↔	↔
Saxagliptin	↑	↑	↑	↑	↑	↔	↓	↓	↓	↔	↔	↔	↔	↔	↔	↔	↑	↑	↔	↔	↔	↔
Sitagliptin	↑ d	↑ d	↑ d	↑ d	↑ d	↔	↓	↓	↓	↔	↔	↔	↔	↔	↔	↔	↑ d	↑ d	↔	↔	↔	↔
Vildagliptin	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔
<b>SGLT-2 inhibitors</b>																						
Canagliflozin	↔	↓	↔	↓	↓	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔ e	↔	↔	↔ e ↔ e
Dapagliflozin	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔
Empagliflozin	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↑	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔
<b>Others</b>																						
Acarbose	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔

**Interactions with CAB/RPV long acting injections**  
Pharmacokinetic interactions shown are mostly with RPV.  
QT interactions shown are with RPV.

**Interactions with Abacavir (ABC), Lamivudine (3TC) or Zidovudine (ZDV)**  
ABC: No clinically relevant interactions expected.  
3TC: No clinically relevant interactions expected.  
ZDV: No clinically relevant interactions expected.

**Interactions with Ibalizumab**  
None

**Colour Legend**

- No clinically significant interaction expected.
- These drugs should not be coadministered.
- Potential interaction which may require a dose adjustment or close monitoring.
- Potential interaction predicted to be of weak intensity.  
No *a priori* dosage adjustment is recommended.

**Text Legend**

- ↑ Potential increased exposure of the anti-diabetic drug
- ↓ Potential decreased exposure of the anti-diabetic drug
- ↔ No significant effect
- ↑ Potential increased exposure of HIV drug
- ↓ Potential decreased exposure of HIV drug

Numbers refer to increase or decrease in AUC as observed in drug-drug interaction studies.

**Notes**

- a Close monitoring is recommended when starting or stopping the combination of these antiretrovirals and metformin as a dose adjustment of metformin may be necessary.
- b Caution is needed when coadministering atazanavir and GLP-1 agonists due to their potential to inhibit gastric secretion (and in some cases to slow gastric emptying), thereby reducing the absorption of atazanavir. Consider taking atazanavir 2-4 hours before the GLP-1 agonist.
- c Caution is needed when coadministering oral rilpivirine and GLP-1 agonists due to their potential to inhibit gastric secretion (and in some cases to slow gastric emptying), thereby reducing the absorption of rilpivirine. Consider taking oral rilpivirine 4 hours before the GLP-1 agonist.
- d Increase in anti-diabetic drug exposure is not considered as clinically significant as the drug is mainly eliminated unchanged and has a large safety window.
- e Caution is recommended when coadministering canagliflozin in the long term with tenofovir-DF due to potential additive bone toxicities.