

Antipsychotic Treatment Selector

Charts revised July 2019. Full information available at www.hiv-druginteractions.org

For personal use only. Not for distribution. For personal use only. Not for distribution. For personal use only. Not for distribution. For personal use only. Not for distribution.

	ATV/c	ATV/r	DRV/c	DRV/r	LPV/r	DOR	EFV	ETV	NVP	RPV	MVC	BIC/F/TAF	DTG	EVG/c/F/TAF	EVG/c/F/TDF	RAL	ABC	FTC or 3TC	F/TAF	TDF	ZDV
Atypical Antipsychotics																					
Amisulpride	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	
Aripiprazole	↑	↑	↑	↑	↑	↔	↓	↓	↓	↔	↔	↔	↔	↑	↑	↑	↔	↔	↔	↔	
Asenapine	↑	↓	↑	↓	↓	↔	↓	↓	↓	↔	↔	↔	↔	↑	↑	↑	↔	↔	↔	↔	
Clozapine	↑♥	↑♥	↑	↑	↑♥	↔	↓	↓	↓	↔♥	↔	↔	↔	↑	↑	↑	↔	↔	↔	↔	
Olanzapine	↔	↓	↔	↓	↓	↔	↓	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	
Paliperidone	↑	↑	↑	↑	↑	↔	↓	↓	↓	↔	↔	↔	↔	↑	↑	↑	↔	↔	↔	↔	
Quetiapine	↑ ^b ♥	↑ ^b ♥	↑ ^b	↑ ^b	↑ ^b ♥	↔	↓	↓	↓	↔♥	↔	↔	↔	↑ ^b	↑ ^b	↔	↔	↔	↔	↔ ^a	
Risperidone	↑	↑	↑	↑	↑	↔	↓	↓	↓	↔	↔	↔	↔	↑	↑	↑	↔	↔	↔	↔	
Zotepine	↑♥	↑♥	↑	↑	↑♥	↔	↓	↓	↓	↔	↔	↔	↔	↑	↑	↑	↔	↔	↔	↔	
Phenothiazines																					
Chlorpromazine	↑♥	↑♥	↑	↑	↑♥	↔	↔	↔	↔♥	↔	↔	↔	↔	↑	↑	↑	↔	↔	↔	↔	
Fluphenazine	↑♥	↑♥	↑	↑	↑♥	↔	↔	↔	↔♥	↔	↔	↔	↔	↑	↑	↑	↔	↔	↔	↔	
Levomepromazine	↑♥	↑♥	↑	↑	↑♥	↔	↔	↔	↔♥	↔	↔	↔	↔	↑	↑	↑	↔	↔	↔	↔	
Perazine	↑♥	↑♥	↑	↑	↑♥	↔	↓	↓	↓	↔♥	↔	↔	↔	↑	↑	↑	↔	↔	↔	↔	
Periciazine	↑♥	↑♥	↑	↑	↑♥	↔	↔	↔	↔♥	↔	↔	↔	↔	↑	↑	↑	↔	↔	↔	↔ ^a	
Perphenazine	↑♥	↑♥	↑	↑	↑♥	↔	↔	↔	↔♥	↔	↔	↔	↔	↑	↑	↑	↔	↔	↔	↔ ^a	
Pimozide	↑♥	↑♥	↑	↑	↑♥	↔	↑ ^c	↓	↓	↔♥	↔	↔	↔	↑	↑	↑	↔	↔	↔	↔	
Prochlorperazine	↑♥	↑♥	↑	↑	↑♥	↔	↔	↔	↔♥	↔	↔	↔	↔	↑	↑	↑	↔	↔	↔	↔ ^a	
Thioridazine	↑♥	↑♥	↑	↑	↑♥	↓	↓	↓	↓	↓♥	↓	↓	↓	↑	↑	↑	↔	↔	↔	↔ ^a	
Others																					
Haloperidol	↑♥	↑♥	↑	↑	↑♥	↔	↓	↓	↓	↔♥	↑	↔	↔	↑	↑	↑	↔	↔	↔	↔	
Iloperidone	↑♥	↑♥	↑	↑	↑♥	↔	↓	↓	↓	↔♥	↔	↔	↔	↑	↑	↑	↔	↔	↔	↔	
Pipotiazine	↑♥	↑♥	↑	↑	↑♥	↔	↔	↔	↔♥	↔	↔	↔	↔	↑	↑	↑	↔	↔	↔	↔ ^a	
Sulpiride	↔♥	↔♥	↔	↔	↔♥	↔	↔	↔	↔♥	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	
Tiapride	↔♥	↔♥	↔	↔	↔♥	↔	↔	↔	↔♥	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	
Ziprasidone	↑♥	↑♥	↑	↑	↑♥	↔	↓	↓	↓	↔♥	↔	↔	↔	↑	↑	↑	↔	↔	↔	↔	
Zuclopentixol	↑♥	↑♥	↑	↑	↑♥	↔	↓	↓	↓	↔♥	↔	↔	↔	↑	↑	↑	↔	↔	↔	↔ ^a	

Colour Legend

- No clinically significant interaction expected.
- These drugs should not be coadministered.
- Potential interaction which may require a dose adjustment or close monitoring.
- Potential interaction predicted to be of weak intensity. No *a priori* dosage adjustment is recommended.

Text Legend

- ↑ Potential increased exposure of the antipsychotic
- ↓ Potential decreased exposure of the antipsychotic
- ↔ No significant effect
- ↑ Potential increased exposure of HIV drug
- ↓ Potential decreased exposure of HIV drug
- ↔ One or both drugs may cause QT and/or PR prolongation. ECG monitoring is advised if coadministered with atazanavir or lopinavir; caution is advised with rilpivirine as supratherapeutic doses of rilpivirine (75 and 300 mg once daily) were shown to prolong the QT interval.

Notes

- a Potential haematological toxicity
- b Coadministration contraindicated in the European SPC however US Prescribing Information recommends quetiapine should be reduced to one sixth of the original dose if coadministered with a potent CYP3A4 inhibitor. The charts reflect the more cautious option.
- c Coadministration contraindicated in the European SPC but not in the US Prescribing Information for Sustiva. The charts reflect the more cautious option.
- d No effect on FTC or TAF is expected, but bictegravir concentrations may decrease.