

# Anti-Hypertensive Treatment Selector

Charts reviewed June 2022. Full information available at [www.hiv-druginteractions.org](http://www.hiv-druginteractions.org)

For personal use only. Not for distribution. For personal use only. Not for distribution. For personal use only. Not for distribution. For personal use only. Not for distribution.

	ATV/c	ATV/r	DRV/c	DRV/r	LPV/r	DOR	EFV	ETV	NVP	RPV oral	FTR	MVC	BIC/F/TAF	CAB oral	CAB/RPV	DTG	EVG/c/F/TAF	EVG/c/F/TDF	RAL	FTC/TAF	FTC/TDF	TDF
<b>ACE Inhibitors</b>																						
Captopril	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔
Cilazapril	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔
Enalapril	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔
Fosinopril	↔	↑	↔	↑	↑	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔
Lisinopril	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔
Perindopril	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔
Quinapril	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔
Ramipril	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔
Trandolapril	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔
<b>Angiotensin Antagonists</b>																						
Candesartan	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔
Eprosartan	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔
Irbesartan	↔	↓	↔	↓	↓	↔	↑	↑	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔
Losartan	↔	↓ a	↔	↓ a	↓ a	↔	↑ b	↑ b	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔
Olmesartan	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔
Telmisartan	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔
Valsartan	↑	↑	↑	↑	↑	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔
<b>Diuretics</b>																						
Amiloride	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔
Bendroflumethiazide	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔
Chlorthalidone	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔
Eplerenone	↑	↑	↑	↑	↑	↔	↓	↓	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔
Furosemide	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔
Hydrochlorothiazide	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔
Indapamide	↑	↑	↑	↑	↑	↔	↓	↓	↓	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔
Torsemide	↔	↓	↔	↓	↓	↔	↑	↑	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔
Triamterene	↔	↓	↔	↓	↓	↔	↑	↑	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔
Xipamide	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔
<b>Others</b>																						
Clonidine	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔
Doxazosin	↑	↑	↑	↑	↑	↔	↓	↓	↓	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔
Hydralazine	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔
Methyldopa	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔
Moxonidine	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔
Prazosin	↑?	↑?	↑?	↑?	↑?	↔	↓?	↓?	↓?	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔
Sacubitril	↑	↑	↑	↑	↑	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔
Spirololactone	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔

**Interactions with CAB/RPV long acting injections**  
Pharmacokinetic interactions shown are mostly with RPV. QT interactions shown are with RPV.

**Interactions with Abacavir (ABC), Lamivudine (3TC) or Zidovudine (ZDV)**  
ABC: no clinically relevant interactions expected.  
3TC: increased amiloride and 3TC exposure when coadministered.  
ZDV: potential additive haematological toxicity with methyldopa.

**Interactions with Ibalizumab**  
None

**Colour Legend**

- No clinically significant interaction expected.
- These drugs should not be coadministered.
- Potential interaction which may require a dose adjustment or close monitoring.
- Potential interaction predicted to be of weak intensity. No *a priori* dosage adjustment is recommended.

**Text Legend**

- ↑ Potential increased exposure of the antihypertensive
- ↓ Potential decreased exposure of the antihypertensive
- ↔ No significant effect
- ↕ One or both drugs may cause QT and/or PR prolongation. ECG monitoring is advised if coadministered with atazanavir or lopinavir. Rilpivirine and fostemsavir were shown to prolong the QT interval at supratherapeutic doses. Caution is advised with rilpivirine. ECG monitoring is advised with fostemsavir and drugs with a known QT prolongation risk.
- Numbers refer to increase or decrease in AUC as observed in drug-drug interaction studies.
- ↑ Potential increased exposure of HIV drug

**Notes**

- a Concentrations of parent drug decreased but concentrations of active metabolite increased.
- b Concentrations of parent drug increased but concentrations of active metabolite decreased.
- c No effect on emtricitabine is expected. Any effect on renal elimination of tenofovir is likely to be limited. No *a priori* dosage adjustment is recommended.
- d Use with caution in patients with a history of postural hypotension or on concomitant medicinal products known to lower blood pressure, and those at increased risk of cardiovascular events.
- e Hydralazine has some nephrotoxic potential. Use of tenofovir-DF should be avoided with concurrent or recent use of a nephrotoxic medicinal product. If coadministration is unavoidable, renal function should be monitored closely.
- f The renal transporter involved in moxonidine excretion is unknown. No interaction is expected with emtricitabine, however, an interaction with tenofovir-DF cannot be excluded.
- g The renal transporter involved in moxonidine excretion is unknown. An interaction with tenofovir-DF cannot be excluded.
- h No interaction is expected with emtricitabine. Tenofovir could potentially compete with LMQ657 renal elimination but the clinical relevance is unclear. Start with the lowest recommended dose of sacubitril and titrate dosage as tolerated by the patient.
- i Tenofovir could potentially compete with LMQ657 renal elimination but the clinical relevance is unclear. Start with the lowest recommended dose of sacubitril and titrate dosage as tolerated by the patient.

Note: although some drug interactions are predicted to potentially require a dosage adjustment based on the drug's metabolic pathway, clinical experience with a particular antihypertensive and HIV drug may indicate that dosage adjustments are not an *a priori* requirement.