

Lipid-Lowering Therapies

Charts revised November 2024. Full information available at www.hiv-druginteractions.org

For personal use only. Not for distribution. For personal use only. Not for distribution. For personal use only. Not for distribution. For personal use only. Not for distribution.

	ATV/c	ATV/r	DRV/c	DRV/r	LPV/r	DOR	EFV	ETV	NVP	RPV oral	FTR	LEN	MVC	BIC/F/TAF	CAB oral	CAB/RPV	DTG	EVG/c/F/TAF	EVG/c/F/TDF	RAL	FTC/TAF	FTC/TDF
Statins																						
Atorvastatin	↑822%	↑	↑290%	↑	↑490%	↓2%	↓43%	↓37%	↓	↑4%	↑	↑	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔
Fluvastatin	↑	↑	↑	↑	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔
Lovastatin	↑	↑	↑	↑	↑	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔
Pitavastatin	↑	↑ ^a	↑	↓26%	↓20%	↔	↓11%	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔
Pravastatin	↑	↑	↑	↑81%	↔	↔	↓44%	↓	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔
Rosuvastatin	↑242%	↑213%	↑93%	↑48%	↑108%	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔
Simvastatin	↑	↑	↑	↑	↑	↔	↓68%	↓	↓	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔
Fibrates																						
Bezafibrate	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔
Clofibrate	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔
Fenofibrate	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔
Gemfibrozil	↔	↓	↔	↓	↓41%	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔
Other																						
Alirocumab	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔
Bempedoic acid	↔	↓	↔	↓	↓	↔	↑	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔
Evolocumab	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔
Ezetimibe	↑	↑	↑	↓↑	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔
Icosapent ethyl	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔
Inclisiran	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔
Lomitapide	↑	↑	↑	↑	↑	↔	↓	↓	↓	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔	↔

Interactions with CAB/RPV long acting injections

Pharmacokinetic interactions shown are mostly with RPV. QT interactions shown are with RPV.

Interactions with Lenacapavir

Residual LEN may affect exposure of sensitive CYP3A4 substrates initiated within 9 months after stopping subcutaneous LEN.

Interactions with Ibalizumab

None

Interactions with Abacavir (ABC), Lamivudine (3TC), Tenofovir-DF (TDF) or Zidovudine (ZDV)

ABC: No clinically relevant interactions expected.

3TC: No clinically relevant interactions expected.

TDF: No clinically relevant interactions expected.

ZDV: Coadministration with bempedoic acid may increase the risk of haematotoxicity. Use with caution and closely monitor haematological parameters.

Colour Legend

- █ No clinically significant interaction expected.
- █ These drugs should not be coadministered.
- █ Potential interaction which may require a dose adjustment or close monitoring.
- █ Potential interaction predicted to be of weak intensity. No *a priori* dosage adjustment is recommended.

Text Legend

- ↑ Potential increased exposure of the lipid-lowering drug
- ↓ Potential decreased exposure of the lipid-lowering drug
- ↔ No significant effect

Numbers refer to increase or decrease in AUC as observed in drug-drug interaction studies.

Notes

- ^a Coadministration with unboosted atazanavir increased pitavastatin AUC by 31%.