









# Interactions with HBV Treatment

**Please note that if a drug is not listed it cannot automatically be assumed it is safe to coadminister.**

ADV, Adefovir; ETV, Entecavir; LAM, Lamivudine; PEG IFN, Peginterferon; RBV, Ribavirin; TBV, Telbivudine; TAF, Tenofovir alafenamide; TDF, Tenofovir-DF.

	ADV	ETV	LAM	PEG IFN alfa-2a	PEG IFN alfa-2b	RBV	TBV	TAF	TDF
<b>HIV Drugs</b>									
<b>Entry/Integrase Inhibitors</b>									
Albuvirtide	◆	◆	◆	◆	◆	◆	◆	◆	◆
Bictegravir/FTC/TAF	●	◆	◆	◆	◆	◆	◆	◆	◆
Cabotegravir (oral)	◆	◆	◆	◆	◆	◆	◆	◆	◆
Cabotegravir/rilpivirine (LA)	◆	◆	◆	◆	◆	◆	◆	◆	◆
Dolutegravir	◆	◆	◆	◆	◆	◆	◆	◆	◆
Dolutegravir/rilpivirine	◆	◆	◆	◆	◆	◆	◆	◆	◆
Elvitegravir/cobi /FTC/TAF	●	◆	◆	◆	◆	◆	◆	◆	◆
Elvitegravir/cobi/FTC/TDF	●	◆	◆	◆	◆	◆	◆	◆	◆
Enfuvirtide	◆	◆	◆	◆	◆	◆	◆	◆	◆
Fostemsavir	◆	◆	◆	◆	◆	◆	◆	◆	◆
Ibalizumab-uyk	◆	◆	◆	◆	◆	◆	◆	◆	◆
Maraviroc	◆	◆	◆	◆	◆	◆	◆	◆	◆
Raltegravir	◆	◆	◆	◆	◆	◆	◆	◆	◆
<b>NNRTIs</b>									
Dapivirine	◆	◆	◆	◆	◆	◆	◆	◆	◆
Dolutegravir/rilpivirine	◆	◆	◆	◆	◆	◆	◆	◆	◆
Doravirine	◆	◆	◆	◆	◆	◆	◆	◆	◆
Doravirine/3TC/TDF	●	◆	◆	◆	◆	◆	◆	◆	◆
Efavirenz	◆	◆	◆	◆	◆	◆	◆	◆	◆
Etravirine	◆	◆	◆	◆	◆	◆	◆	◆	◆
Nevirapine	◆	◆	◆	◆	◆	◆	◆	◆	◆
Rilpivirine	◆	◆	◆	◆	◆	◆	◆	◆	◆
Rilpivirine/FTC/TAF	●	◆	◆	◆	◆	◆	◆	◆	◆
<b>NRTIs</b>									
Abacavir	◆	◆	◆	◆	◆	◆	◆	◆	◆
Didanosine	◆	◆	◆	◆	◆	◆	◆	◆	◆
Emtricitabine (FTC)	◆	◆	◆	◆	◆	◆	◆	◆	◆
Emtricitabine + TAF	●	◆	◆	◆	◆	◆	◆	◆	◆
Emtricitabine + TDF	●	◆	◆	◆	◆	◆	◆	◆	◆
Lamivudine	◆	◆	◆	◆	◆	◆	◆	◆	◆
Stavudine	◆	◆	◆	◆	◆	◆	◆	◆	◆
Tenofovir-DF	●	◆	◆	◆	◆	◆	◆	◆	◆
Zidovudine	◆	◆	◆	◆	◆	◆	◆	◆	◆
<b>Protease Inhibitors</b>									
Atazanavir alone	◆	◆	◆	◆	◆	◆	◆	◆	◆
Atazanavir/cobicistat	◆	◆	◆	◆	◆	◆	◆	◆	◆
Atazanavir + ritonavir	◆	◆	◆	◆	◆	◆	◆	◆	◆
Darunavir/cobicistat	◆	◆	◆	◆	◆	◆	◆	◆	◆
Darunavir/cobi/FTC/TAF	●	◆	◆	◆	◆	◆	◆	◆	◆
Darunavir + ritonavir	◆	◆	◆	◆	◆	◆	◆	◆	◆
Fosamprenavir	◆	◆	◆	◆	◆	◆	◆	◆	◆
Indinavir	◆	◆	◆	◆	◆	◆	◆	◆	◆
Lopinavir	◆	◆	◆	◆	◆	◆	◆	◆	◆
Ritonavir	◆	◆	◆	◆	◆	◆	◆	◆	◆
Tipranavir	◆	◆	◆	◆	◆	◆	◆	◆	◆

	ADV	ETV	LAM	PEG IFN alfa-2a	PEG IFN alfa-2b	RBV	TBV	TAF	TDF
<b>Hypertension/Heart Failure Agents</b>									
Acebutolol	◆	◆	◆	◆	◆	◆	◆	◆	◆
Aliskiren	◆	◆	◆	◆	◆	◆	◆	◆	◆
Ambrisentan	◆	◆	◆	◆	◆	◆	◆	◆	◆
Amiloride	◆	◆	◆	◆	◆	◆	◆	◆	◆
Azilsartan	◆	◆	◆	◆	◆	◆	◆	◆	◆
Benazepril	◆	◆	◆	◆	◆	◆	◆	◆	◆
Bendroflumethiazide	◆	◆	◆	◆	◆	◆	◆	◆	◆
Bosentan	◆	◆	◆	◆	◆	◆	◆	◆	◆
Bumetanide	◆	◆	◆	◆	◆	◆	◆	◆	◆
Candesartan	◆	◆	◆	◆	◆	◆	◆	◆	◆
Captopril	◆	◆	◆	◆	◆	◆	◆	◆	◆
Chlorthalidone	◆	◆	◆	◆	◆	◆	◆	◆	◆
Cilazapril	◆	◆	◆	◆	◆	◆	◆	◆	◆
Clevidipine	◆	◆	◆	◆	◆	◆	◆	◆	◆
Clonidine	◆	◆	◆	◆	◆	◆	◆	◆	◆
Doxazosin	◆	◆	◆	◆	◆	◆	◆	◆	◆
Enalapril	◆	◆	◆	◆	◆	◆	◆	◆	◆
Eplerenone	◆	◆	◆	◆	◆	◆	◆	◆	◆
Epoprostenol	◆	◆	◆	◆	◆	◆	◆	◆	◆
Eprosartan	◆	◆	◆	◆	◆	◆	◆	◆	◆
Fosinopril	◆	◆	◆	◆	◆	◆	◆	◆	◆
Furosemide	◆	◆	◆	◆	◆	◆	◆	◆	◆
Hydralazine	◆	◆	◆	◆	◆	◆	◆	◆	◆
Hydrochlorothiazide	◆	◆	◆	◆	◆	◆	◆	◆	◆
Iloprost	◆	◆	◆	◆	◆	◆	◆	◆	◆
Indapamide	◆	◆	◆	◆	◆	◆	◆	◆	◆
Irbesartan	◆	◆	◆	◆	◆	◆	◆	◆	◆
Isradipine	◆	◆	◆	◆	◆	◆	◆	◆	◆
Ivabradine	◆	◆	◆	◆	◆	◆	◆	◆	◆
Lacidipine	◆	◆	◆	◆	◆	◆	◆	◆	◆
Lercanidipine	◆	◆	◆	◆	◆	◆	◆	◆	◆
Lisinopril	◆	◆	◆	◆	◆	◆	◆	◆	◆
Losartan	◆	◆	◆	◆	◆	◆	◆	◆	◆
Macitentan	◆	◆	◆	◆	◆	◆	◆	◆	◆
Methyldopa	◆	◆	◆	◆	◆	◆	◆	◆	◆
Metolazone	◆	◆	◆	◆	◆	◆	◆	◆	◆
Moxonidine	◆	◆	◆	◆	◆	◆	◆	◆	◆
Olmesartan	◆	◆	◆	◆	◆	◆	◆	◆	◆
Perindopril	◆	◆	◆	◆	◆	◆	◆	◆	◆
Prazosin	◆	◆	◆	◆	◆	◆	◆	◆	◆
Quinapril	◆	◆	◆	◆	◆	◆	◆	◆	◆
Ramipril	◆	◆	◆	◆	◆	◆	◆	◆	◆
Ranolazine	◆	◆	◆	◆	◆	◆	◆	◆	◆
Rilmendidine	◆	◆	◆	◆	◆	◆	◆	◆	◆
Riociguat	◆	◆	◆	◆	◆	◆	◆	◆	◆
Sacubitril/valsartan	◆	◆	◆	◆	◆	◆	◆	◆	◆
Selexipag	◆	◆	◆	◆	◆	◆	◆	◆	◆
Sildenafil	◆	◆	◆	◆	◆	◆	◆	◆	◆
Spirolactone	◆	◆	◆	◆	◆	◆	◆	◆	◆
Tadalafil	◆	◆	◆	◆	◆	◆	◆	◆	◆
Telmisartan	◆	◆	◆	◆	◆	◆	◆	◆	◆
Torsemide	◆	◆	◆	◆	◆	◆	◆	◆	◆
Trandolapril	◆	◆	◆	◆	◆	◆	◆	◆	◆
Treprostinil	◆	◆	◆	◆	◆	◆	◆	◆	◆
Valsartan	◆	◆	◆	◆	◆	◆	◆	◆	◆
Xipamide	◆	◆	◆	◆	◆	◆	◆	◆	◆
Zofenopril	◆	◆	◆	◆	◆	◆	◆	◆	◆

For personal use only. Not for distribution.

**Key to symbols**

●	These drugs should not be coadministered
■	Potential clinically significant interaction that is likely to require additional monitoring, alteration of drug dosage or timing of administration
▲	Potential interaction likely to be of weak intensity. Additional action/monitoring or dosage adjustment is unlikely to be required
◆	No clinically significant interaction expected

**Notes**

- Further information is available at [www.hep-druginteractions.org](http://www.hep-druginteractions.org)
- Predicted interactions are based on known metabolic pathways and routes of clearance.
- Caution is required in patients with hepatic impairment as this may also increase drug levels and require dose modification.
- Where advice differs between countries, the charts reflect the more cautious option.

© Liverpool Drug Interactions Group,  
 Liverpool Drug Interactions Group, University of Liverpool, 3rd Floor William Henry Duncan Building, 6 West Derby Street, Liverpool, L7 8TX.  
 We aim to ensure that information is accurate and consistent with current knowledge and practice. However, the University of Liverpool and its servants or agents shall not be responsible or in any way liable for the continued currency of information in this publication whether arising from negligence or otherwise howsoever or for any consequences arising therefrom. The University of Liverpool expressly exclude liability for errors, omissions or inaccuracies to the fullest extent permitted by law.

# Interactions with HBV Treatment

Charts revised January 2023. Full information available at [www.hep-druginteractions.org](http://www.hep-druginteractions.org)

Page 6 of 6

**Please note that if a drug is not listed it cannot automatically be assumed it is safe to coadminister.**

ADV, Adefovir; ETV, Entecavir; LAM, Lamivudine; PEG IFN, Peginterferon; RBV, Ribavirin; TBV, Telbivudine; TAF, Tenofovir alafenamide; TDF, Tenofovir-DF.

	ADV	ETV	LAM	PEG IFN alfa-2a	PEG IFN alfa-2b	RBV	TBV	TAF	TDF
<b>Illicit/Recreational</b>									
Alcohol	◆	◆	◆	◆	◆	◆	◆	◆	◆
Amphetamine	◆	◆	◆	◆	◆	◆	◆	◆	◆
Cannabis	◆	◆	◆	◆	◆	◆	◆	◆	◆
Carfentanil	◆	◆	◆	◆	◆	◆	◆	◆	◆
Cocaine	◆	◆	◆	◆	◆	◆	◆	◆	◆
Ecstasy (MDMA)	◆	◆	◆	◆	◆	◆	◆	◆	◆
Fentanyl (Prescribed)	◆	◆	◆	◆	◆	◆	◆	◆	◆
GHB (Gamma-hydroxybutyrate)	◆	◆	◆	◆	◆	◆	◆	◆	◆
Heroin	◆	◆	◆	◆	◆	◆	◆	◆	◆
LSD (Lysergic acid diethylamide)	◆	◆	◆	◆	◆	◆	◆	◆	◆
Mephedrone	◆	◆	◆	◆	◆	◆	◆	◆	◆
Methamphetamine	◆	◆	◆	◆	◆	◆	◆	◆	◆
Phencyclidine (PCP)	◆	◆	◆	◆	◆	◆	◆	◆	◆
<b>Immunosuppressants</b>									
Adalimumab	◆	◆	◆	▲	▲	■	◆	◆	◆
Alemtuzumab	◆	◆	◆	■	■	■	◆	◆	◆
Anakinra	◆	◆	◆	■	■	■	◆	◆	◆
Azathioprine	◆	◆	◆	■	■	■	◆	◆	◆
Baricitinib	◆	◆	◆	■	■	■	◆	◆	◆
Basiliximab	◆	◆	◆	◆	◆	◆	◆	◆	◆
Belimumab	◆	◆	◆	◆	◆	◆	◆	◆	◆
Brodalumab	◆	◆	◆	◆	◆	◆	◆	◆	◆
Canakinumab	◆	◆	◆	■	■	■	◆	◆	◆
Ciclosporin	◆	◆	◆	◆	◆	◆	■	■	■
Eculizumab	◆	◆	◆	◆	◆	◆	◆	◆	◆
Etanercept	◆	◆	◆	◆	◆	◆	◆	◆	◆
Fingolimod	◆	◆	◆	◆	◆	◆	◆	◆	◆
Golimumab	◆	◆	◆	■	■	■	◆	◆	◆
Guselkumab	◆	◆	◆	◆	◆	◆	◆	◆	◆
Infliximab	◆	◆	◆	◆	◆	◆	◆	◆	◆
Lenalidomide	◆	◆	◆	◆	◆	◆	◆	◆	◆
Mycophenolate	◆	◆	◆	■	■	■	◆	◆	■
Pirfenidone	◆	◆	◆	◆	◆	◆	◆	◆	◆
Ravulizumab	◆	◆	◆	◆	◆	◆	◆	◆	◆
Risankizumab	◆	◆	◆	◆	◆	◆	◆	◆	◆
Sarilumab	◆	◆	◆	■	■	■	◆	◆	◆
Secukinumab	◆	◆	◆	◆	◆	◆	◆	◆	◆
Siltuximab	◆	◆	◆	■	■	■	◆	◆	◆
Sirolimus	◆	◆	◆	◆	◆	◆	◆	◆	▲
Tacrolimus	◆	◆	◆	◆	◆	◆	◆	◆	▲
Tildrakizumab	◆	◆	◆	◆	◆	◆	◆	◆	◆
Tocilizumab	◆	◆	◆	■	■	■	◆	◆	◆
Tralokinumab	◆	◆	◆	◆	◆	◆	◆	◆	◆
Upadacitinib	◆	◆	◆	■	■	■	◆	◆	◆
Ustekinumab	◆	◆	◆	◆	◆	◆	◆	◆	◆
Vedolizumab	◆	◆	◆	◆	◆	◆	◆	◆	◆
<b>Lipid Lowering Agents</b>									
Alirocumab	◆	◆	◆	◆	◆	◆	◆	◆	◆
Atorvastatin	◆	◆	◆	◆	◆	◆	■	◆	◆
Bezafibrate	◆	◆	◆	◆	◆	◆	◆	◆	◆
Evolocumab	◆	◆	◆	◆	◆	◆	◆	◆	◆
Ezetimibe	◆	◆	◆	◆	◆	◆	◆	◆	◆
Fenofibrate	◆	◆	◆	◆	◆	◆	■	◆	◆
Fish oils	◆	◆	◆	◆	◆	◆	◆	◆	◆
Fluvastatin	◆	◆	◆	◆	◆	◆	◆	◆	◆
Gemfibrozil	◆	◆	◆	◆	◆	◆	■	◆	◆
Lovastatin	◆	◆	◆	◆	◆	◆	◆	◆	◆
Pitavastatin	◆	◆	◆	◆	◆	◆	■	◆	◆
Pravastatin	◆	◆	◆	◆	◆	◆	◆	◆	◆
Rosuvastatin	◆	◆	◆	◆	◆	◆	■	◆	◆
Simvastatin	◆	◆	◆	◆	◆	◆	■	◆	◆

	ADV	ETV	LAM	PEG IFN alfa-2a	PEG IFN alfa-2b	RBV	TBV	TAF	TDF
<b>Other Drugs</b>									
Acamprosate	◆	◆	◆	◆	◆	◆	◆	◆	◆
Acetazolamide	■	■	◆	◆	◆	◆	◆	◆	■
Acitretin	◆	◆	◆	◆	◆	◆	◆	◆	◆
Activated charcoal	▲	▲	▲	◆	◆	▲	▲	▲	▲
Allopurinol	◆	◆	◆	■	■	■	◆	◆	◆
Atomoxetine	◆	◆	◆	◆	◆	◆	◆	◆	◆
Atropine	◆	◆	◆	◆	◆	◆	◆	◆	◆
Baclofen	◆	◆	◆	◆	◆	◆	◆	◆	◆
Bamlanivimab	◆	◆	◆	◆	◆	◆	◆	◆	◆
Benralizumab	◆	◆	◆	◆	◆	◆	◆	◆	◆
Bethahistine	◆	◆	◆	◆	◆	◆	◆	◆	◆
Bimatoprost	◆	◆	◆	◆	◆	◆	◆	◆	◆
Biperiden	◆	◆	◆	◆	◆	◆	◆	◆	◆
Brinzolamide	◆	◆	◆	◆	◆	◆	◆	◆	◆
Brolucizumab	◆	◆	◆	◆	◆	◆	◆	◆	◆
Bromocriptine	◆	◆	◆	◆	◆	◆	◆	◆	◆
Calcitonin	◆	◆	◆	◆	◆	◆	◆	◆	◆
Calcium carbimide	◆	◆	◆	◆	◆	◆	◆	◆	◆
Calcium resonium	◆	◆	◆	◆	◆	◆	◆	◆	◆
Cannabidiol	◆	◆	◆	◆	◆	◆	◆	◆	◆
Carbimazole	◆	◆	◆	◆	◆	◆	◆	◆	◆
Carisoprodol	◆	◆	◆	◆	◆	◆	◆	◆	◆
Casirivimab/imdevimab	◆	◆	◆	◆	◆	◆	◆	◆	◆
Cilostazol	◆	◆	◆	◆	◆	◆	◆	◆	◆
Clomifene	◆	◆	◆	◆	◆	◆	◆	◆	◆
Colchicine	◆	◆	◆	◆	◆	◆	◆	◆	◆
Colestyramine	▲	▲	▲	◆	◆	▲	▲	▲	▲
Conivaptan	◆	◆	◆	◆	◆	◆	◆	◆	◆
Convalescent plasma (COVID-19)	◆	◆	◆	◆	◆	◆	◆	◆	◆
COVID-19 vaccinations	◆	◆	◆	◆	◆	◆	◆	◆	◆
Crizanlizumab	◆	◆	◆	◆	◆	◆	◆	◆	◆
Cyclobenzaprine	◆	◆	◆	◆	◆	◆	◆	◆	◆
Cytisine	◆	◆	◆	◆	◆	◆	◆	◆	◆
Darbepoetin	◆	◆	◆	◆	◆	◆	◆	◆	◆
Deferiprone	◆	◆	◆	◆	◆	◆	◆	◆	◆
Denosumab	◆	◆	◆	◆	◆	◆	◆	◆	◆
Dexamfetamine	◆	◆	◆	◆	◆	◆	◆	◆	◆
Dextromethorphan	◆	◆	◆	◆	◆	◆	◆	◆	◆
Disulfiram	◆	◆	◆	■	■	◆	◆	◆	◆
Donepezil	◆	◆	◆	◆	◆	◆	◆	◆	◆
Dorzolamide	◆	◆	◆	◆	◆	◆	◆	◆	◆
Dupilumab	◆	◆	◆	◆	◆	◆	◆	◆	◆
Eliquis	◆	◆	◆	◆	◆	◆	◆	◆	◆
Emicizumab	◆	◆	◆	◆	◆	◆	◆	◆	◆
Epoetin alfa	◆	◆	◆	◆	◆	◆	◆	◆	◆
Etelcalcetide	◆	◆	◆	◆	◆	◆	◆	◆	◆
Febuxostat	◆	◆	◆	◆	◆	◆	◆	◆	◆
Filgrastim	◆	◆	◆	◆	◆	◆	◆	◆	◆
Flibanserin	◆	◆	◆	◆	◆	◆	◆	◆	◆
Gadopentetate (gadolinium)	◆	◆	◆	◆	◆	◆	◆	◆	◆
Glycerol phenylbutyrate	◆	◆	◆	◆	◆	◆	◆	◆	◆
Goserelin	◆	◆	◆	◆	◆	◆	◆	◆	◆
Idarucizumab	◆	◆	◆	◆	◆	◆	◆	◆	◆
Influenza vaccine	◆	◆	◆	◆	◆	◆	◆	◆	◆
Interferon beta	◆	◆	◆	●	●	■	◆	◆	◆

For personal use only. Not for distribution.

**Key to symbols**

●	These drugs should not be coadministered
■	Potential clinically significant interaction that is likely to require additional monitoring, alteration of drug dosage or timing of administration
▲	Potential interaction likely to be of weak intensity. Additional action/monitoring or dosage adjustment is unlikely to be required
◆	No clinically significant interaction expected

**Notes**

- Further information is available at [www.hep-druginteractions.org](http://www.hep-druginteractions.org)
- Predicted interactions are based on known metabolic pathways and routes of clearance.
- Caution is required in patients with hepatic impairment as this may also increase drug levels and require dose modification.
- Where advice differs between countries, the charts reflect the more cautious option.

© Liverpool Drug Interactions Group,  
 Liverpool Drug Interactions Group, University of Liverpool, 3rd Floor William Henry Duncan Building, 6 West Derby Street, Liverpool, L7 8TX.  
 We aim to ensure that information is accurate and consistent with current knowledge and practice. However, the University of Liverpool and its servants or agents shall not be responsible or in any way liable for the continued currency of information in this publication whether arising from negligence or otherwise howsoever or for any consequences arising therefrom. The University of Liverpool expressly exclude liability for errors, omissions or inaccuracies to the fullest extent permitted by law.

# Interactions with HBV Treatment

Charts revised January 2023. Full information available at [www.hep-druginteractions.org](http://www.hep-druginteractions.org)

**Please note that if a drug is not listed it cannot automatically be assumed it is safe to coadminister.**

ADV, Adefovir; ETV, Entecavir; LAM, Lamivudine; PEG IFN, Peginterferon; RBV, Ribavirin; TBV, Telbivudine; TAF, Tenofovir alafenamide; TDF, Tenofovir-DF.

	ADV	ETV	LAM	PEG IFN alpha-2a	PEG IFN alpha-2b	RBV	TBV	TAF	TDF
<b>Other Drugs Continued</b>									
Isosorbide mononitrate	◆	◆	◆	◆	◆	◆	◆	◆	◆
Isotretinoin	◆	◆	◆	◆	◆	◆	◆	◆	◆
Lanadelumab	◆	◆	◆	◆	◆	◆	◆	◆	◆
Lanreotide	◆	◆	◆	◆	◆	◆	◆	◆	◆
Levothyroxine	◆	◆	◆	◆	◆	◆	◆	◆	◆
Lofexidine	◆	◆	◆	◆	◆	◆	◆	◆	◆
Lumacaftor/ivacaftor	◆	◆	◆	◆	◆	◆	◆	◆	◆
Magnesium	◆	◆	◆	◆	◆	◆	◆	◆	◆
Melatonin	◆	◆	◆	◆	◆	◆	◆	◆	◆
Memantine	◆	◆	◆	◆	◆	◆	◆	◆	◆
Mepolizumab	◆	◆	◆	◆	◆	◆	◆	◆	◆
Methylphenidate	◆	◆	◆	◆	◆	◆	◆	◆	◆
Minoxidil	◆	◆	◆	◆	◆	◆	◆	◆	◆
Modafinil	◆	◆	◆	◆	◆	◆	◆	◆	◆
Naftidrofuryl	◆	◆	◆	◆	◆	◆	◆	◆	◆
Nalmefene	◆	◆	◆	◆	◆	◆	◆	◆	◆
Naloxone	◆	◆	◆	◆	◆	◆	◆	◆	◆
Naltrexone	◆	◆	◆	◆	◆	◆	◆	◆	◆
Neostigmine	◆	◆	◆	◆	◆	◆	◆	◆	◆
Nicorandil	◆	◆	◆	◆	◆	◆	◆	◆	◆
Nusinersen	◆	◆	◆	◆	◆	◆	◆	◆	◆
Ocrelizumab	◆	◆	◆	◆	◆	◆	◆	◆	◆
Orlistat	▲	▲	▲	◆	◆	▲	▲	▲	▲
Penicillamine	◆	◆	◆	◆	◆	◆	◆	◆	◆
Pentoxifylline	◆	◆	◆	◆	◆	◆	◆	◆	◆
Phenylephrine	◆	◆	◆	◆	◆	◆	◆	◆	◆
Pilocarpine	◆	◆	◆	◆	◆	◆	◆	◆	◆
Piracetam	◆	◆	◆	◆	◆	◆	◆	◆	◆
Potassium	◆	◆	◆	◆	◆	◆	◆	◆	◆
Protamine sulphate	◆	◆	◆	◆	◆	◆	◆	◆	◆
Pseudoephedrine	◆	◆	◆	◆	◆	◆	◆	◆	◆
Pyridostigmine	▲	▲	▲	◆	◆	◆	◆	◆	◆
Raloxifene	◆	◆	◆	◆	◆	◆	◆	◆	◆
Ranibizumab	◆	◆	◆	◆	◆	◆	◆	◆	◆
Romosozumab	◆	◆	◆	◆	◆	◆	◆	◆	◆
Sevelamer	▲	▲	▲	◆	◆	▲	▲	▲	▲
Strontium ranelate	◆	◆	◆	◆	◆	◆	◆	◆	◆
Thalidomide	◆	◆	◆	◆	◆	◆	◆	◆	◆
Tranexamic acid	◆	◆	◆	◆	◆	◆	◆	◆	◆
Varenicline	◆	◆	◆	◆	◆	◆	◆	◆	◆

	ADV	ETV	LAM	PEG IFN alpha-2a	PEG IFN alpha-2b	RBV	TBV	TAF	TDF
<b>Oxytocics</b>									
Ergometrine (ergonovine)	◆	◆	◆	◆	◆	◆	◆	◆	◆
Mifepristone	◆	◆	◆	◆	◆	◆	◆	◆	◆
Misoprostol	◆	◆	◆	◆	◆	◆	◆	◆	◆
<b>Parkinsonism Agents</b>									
Benzotropine	◆	◆	◆	◆	◆	◆	◆	◆	◆
Carbidopa	◆	◆	◆	◆	◆	◆	◆	◆	◆
Orphenadrine	◆	◆	◆	◆	◆	◆	◆	◆	◆
Pramipexole	◆	◆	▲	◆	◆	◆	◆	◆	◆
Procyclidine	◆	◆	◆	◆	◆	◆	◆	◆	◆
Rasagiline	◆	◆	◆	◆	◆	◆	◆	◆	◆
Ropinirole	◆	◆	◆	◆	◆	◆	◆	◆	◆
<b>PBC Agents</b>									
Obeticholic acid	▲	◆	◆	◆	◆	◆	◆	◆	◆
Ursodeoxycholic acid	◆	◆	◆	◆	◆	◆	◆	◆	◆
<b>Steroids</b>									
Beclomethasone	◆	◆	◆	◆	◆	◆	◆	◆	◆
Betamethasone	◆	◆	◆	◆	◆	◆	◆	◆	◆
Budesonide	◆	◆	◆	◆	◆	◆	◆	◆	◆
Ciclesonide	◆	◆	◆	◆	◆	◆	◆	◆	◆
Clobetasol (topical)	◆	◆	◆	◆	◆	◆	◆	◆	◆
Clobetasone (topical)	◆	◆	◆	◆	◆	◆	◆	◆	◆
Dexamethasone	◆	◆	◆	◆	◆	◆	◆	◆	◆
Fludrocortisone	◆	◆	◆	◆	◆	◆	◆	◆	◆
Flunisolide	◆	◆	◆	◆	◆	◆	◆	◆	◆
Fluticasone	◆	◆	◆	◆	◆	◆	◆	◆	◆
Hydrocortisone (topical)	◆	◆	◆	◆	◆	◆	◆	◆	◆
Methylprednisolone	◆	◆	◆	◆	◆	◆	◆	◆	◆
Mometasone	◆	◆	◆	◆	◆	◆	◆	◆	◆
Prednicarbate	◆	◆	◆	◆	◆	◆	◆	◆	◆
Prednisone	◆	◆	◆	◆	◆	◆	◆	◆	◆
Triamcinolone	◆	◆	◆	◆	◆	◆	◆	◆	◆
<b>Urological Agents</b>									
Alfuzosin	◆	◆	◆	◆	◆	◆	◆	◆	◆
Desmopressin	◆	◆	◆	◆	◆	◆	◆	◆	◆
Dutasteride	◆	◆	◆	◆	◆	◆	◆	◆	◆
Finasteride	◆	◆	◆	◆	◆	◆	◆	◆	◆
Mirabegron	◆	◆	◆	◆	◆	◆	◆	◆	◆
Silodosin	◆	◆	◆	◆	◆	◆	◆	◆	◆
Solifenacin	◆	◆	◆	◆	◆	◆	◆	◆	◆
Tamsulosin	◆	◆	◆	◆	◆	◆	◆	◆	◆
Tolterodine	◆	◆	◆	◆	◆	◆	◆	◆	◆

For personal use only. Not for distribution.

**Key to symbols**

●	These drugs should not be coadministered
■	Potential clinically significant interaction that is likely to require additional monitoring, alteration of drug dosage or timing of administration
▲	Potential interaction likely to be of weak intensity. Additional action/monitoring or dosage adjustment is unlikely to be required
◆	No clinically significant interaction expected

**Notes**

- Further information is available at [www.hep-druginteractions.org](http://www.hep-druginteractions.org)
- Predicted interactions are based on known metabolic pathways and routes of clearance.
- Caution is required in patients with hepatic impairment as this may also increase drug levels and require dose modification.
- Where advice differs between countries, the charts reflect the more cautious option.

© Liverpool Drug Interactions Group,  
 Liverpool Drug Interactions Group, University of Liverpool, 3rd Floor William Henry Duncan Building, 6 West Derby Street, Liverpool, L7 8TX.  
 We aim to ensure that information is accurate and consistent with current knowledge and practice. However, the University of Liverpool and its servants or agents shall not be responsible or in any way liable for the continued currency of information in this publication whether arising from negligence or otherwise howsoever or for any consequences arising therefrom. The University of Liverpool expressly exclude liability for errors, omissions or inaccuracies to the fullest extent permitted by law.